A0040

Leader in Biomolecular Solutions for Life Science

MET Rabbit pAb

Catalog No.: A0040 5 Publications

Basic Information

Observed MW

140kDa/170kDa

Calculated MW

Polyclonal Antibody

Cross-Reactivity

Applications WB, ELISA

156kDa

Category

Human,Rat



Background

This gene encodes a member of the receptor tyrosine kinase family of proteins and the product of the proto-oncogene MET. The encoded preproprotein is proteolytically processed to generate alpha and beta subunits that are linked via disulfide bonds to form the mature receptor. Further processing of the beta subunit results in the formation of the M10 peptide, which has been shown to reduce lung fibrosis. Binding of its ligand, hepatocyte growth factor, induces dimerization and activation of the receptor, which plays a role in cellular survival, embryogenesis, and cellular migration and invasion. Mutations in this gene are associated with papillary renal cell carcinoma, hepatocellular carcinoma, and various head and neck cancers. Amplification and overexpression of this gene are also associated with multiple human cancers.

Recommended Dilutions

Immunogen Information

WB

1:100 - 1:500

Gene ID 4233

Swiss Prot P08581

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 960-1161 of human MET (NP_000236.2).

Synonyms

DA11; HGFR; AUTS9; RCCP2; c-Met; DFNB97; MET

Product Information

€ www.abclonal.com Source


Isotype lgG

Purification Affinity purification

Storage

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.



Western blot analysis of Rat kidney, using MET antibody (A0040) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 60s.