

A0249

Leader in Biomolecular Solutions for Life Science



## Lamin A/C Rabbit pAb

Catalog No.: A0249

39 Publications

### Basic Information

**Observed MW**

68kDa/72kDa

**Calculated MW**

74kDa

**Category**

Mouse Monoclonal Antibody

**Applications**

WB, IHC-P, IF/ICC, ChIP, ELISA

**Cross-Reactivity**

Human, Mouse, Rat

### Background

The protein encoded by this gene is part of the nuclear lamina, a two-dimensional matrix of proteins located next to the inner nuclear membrane. The lamin family of proteins make up the matrix and are highly conserved in evolution. During mitosis, the lamina matrix is reversibly disassembled as the lamin proteins are phosphorylated. Lamin proteins are thought to be involved in nuclear stability, chromatin structure and gene expression. Vertebrate lamins consist of two types, A and B. Alternative splicing results in multiple transcript variants. Mutations in this gene lead to several diseases: Emery-Dreifuss muscular dystrophy, familial partial lipodystrophy, limb girdle muscular dystrophy, dilated cardiomyopathy, Charcot-Marie-Tooth disease, and Hutchinson-Gilford progeria syndrome.

### Recommended Dilutions

|        |                |
|--------|----------------|
| WB     | 1:500 - 1:1000 |
| IHC-P  | 1:50 - 1:200   |
| IF/ICC | 1:50 - 1:200   |
| ChIP   | 1:20 - 1:100   |

### Immunogen Information

**Gene ID**

4000

**Swiss Prot**

P02545

**Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 403-572 of human Lamin A/C (NP\_733821.1).

**Synonyms**

FPL; IDC; LFP; CDDC; EMD2; FPLD; HGPS; LDP1; LMN1; LMNC; MADA; PRO1; CDCD1; CMD1A; FPLD2; LMNL1; CMT2B1; LGMD1B; Lamin A/C

### Contact

[www.abclonal.com](http://www.abclonal.com)

### Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


## Validation Data




Western blot analysis of various lysates, using Lamin A/C Rabbit pAb (A0249) at 1:1000 dilution.  
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.  
 Lysates/proteins: 25µg per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 1s.




Immunohistochemistry analysis of Lamin A/C in paraffin-embedded mouse brain using Lamin A/C Rabbit pAb (A0249) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of Lamin A/C in paraffin-embedded rat brain using Lamin A/C Rabbit pAb (A0249) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of Lamin A/C in paraffin-embedded rat liver using Lamin A/C Rabbit pAb (A0249) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of NIH/3T3 cells using Lamin A/C Rabbit pAb (A0249) at dilution of 100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (A5007) at 1:500 dilution. Blue: DAPI for nuclear staining.