Leader in Biomolecular Solutions for Life Science

# **PCNA Rabbit pAb**

Catalog No.: A0264 93 Publications



# **Basic Information**

# **Observed MW**

34kDa

### **Calculated MW**

29kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,IHC-P,ELISA

# **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene is found in the nucleus and is a cofactor of DNA polymerase delta. The encoded protein acts as a homotrimer and helps increase the processivity of leading strand synthesis during DNA replication. In response to DNA damage, this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of this gene have been described on chromosome 4 and on the X chromosome.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

IHC-P 1:50 - 1:200

# **Immunogen Information**

**Gene ID Swiss Prot** 5111 P12004

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 162-261 of human PCNA (NP 002583.1).

### **Synonyms**

ATLD2; PCNA

### **Contact**

www.abclonal.com


# **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of extracts of various cell lines, using PCNA antibody (A0264) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).


Exposure time: 3s.

Western blot analysis of extracts of various cell lines, using PCNA antibody (A0264) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).






Immunohistochemistry analysis of paraffinembedded mouse colon using PCNA Rabbit pAb (A0264) at dilution of 1:200 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffinembedded mouse testis using PCNA Rabbit pAb (A0264) at dilution of 1:200 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffinembedded rat testis using PCNA Rabbit pAb (A0264) at dilution of 1:200 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.