

A0270

Leader in Biomolecular Solutions for Life Science



PPAR γ Rabbit pAb

Catalog No.: A0270

63 Publications

Basic Information

Observed MW

57kDa

Calculated MW

58kDa

Category

Antibody

Applications

IHC-P,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

This gene encodes a member of the peroxisome proliferator-activated receptor (PPAR) subfamily of nuclear receptors. PPARs form heterodimers with retinoid X receptors (RXRs) and these heterodimers regulate transcription of various genes. Three subtypes of PPARs are known: PPAR-alpha, PPAR-delta, and PPAR-gamma. The protein encoded by this gene is PPAR-gamma and is a regulator of adipocyte differentiation. Additionally, PPAR-gamma has been implicated in the pathology of numerous diseases including obesity, diabetes, atherosclerosis and cancer. Alternatively spliced transcript variants that encode different isoforms have been described.

Recommended Dilutions

IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

5468

Swiss Prot

P37231

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 100-200 of human PPAR γ (NP_056953.2).

Synonyms

GLM1; CIMT1; NR1C3; PPARG1; PPARG2; PPARG5; PPARGgamma; PPAR γ

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.


Validation Data



Immunohistochemistry analysis of PPAR γ in paraffin-embedded mouse heart using PPAR γ Rabbit pAb (A0270) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of PPAR γ in paraffin-embedded mouse liver using PPAR γ Rabbit pAb (A0270) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of U2OS cells using PPAR γ Rabbit pAb (A0270) at dilution of 1:200 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.