

# DDX3X Mouse mAb

Catalog No.: A10099 **1 Publications**

## Basic Information

**Observed MW**

73kDa

**Calculated MW**

73kDa

**Category**

Primary antibody

**Applications**

ELISA, WB, IF/ICC, IP

**Cross-Reactivity**

Human, Mouse, Rat, Monkey

## Background

The protein encoded by this gene is a member of the large DEAD-box protein family, that is defined by the presence of the conserved Asp-Glu-Ala-Asp (DEAD) motif, and has ATP-dependent RNA helicase activity. This protein has been reported to display a high level of RNA-independent ATPase activity, and unlike most DEAD-box helicases, the ATPase activity is thought to be stimulated by both RNA and DNA. This protein has multiple conserved domains and is thought to play roles in both the nucleus and cytoplasm. Nuclear roles include transcriptional regulation, mRNP assembly, pre-mRNA splicing, and mRNA export. In the cytoplasm, this protein is thought to be involved in translation, cellular signaling, and viral replication. Misregulation of this gene has been implicated in tumorigenesis. This gene has a paralog located in the nonrecombining region of the Y chromosome. Pseudogenes sharing similarity to both this gene and the DDX3Y paralog are found on chromosome 4 and the X chromosome. Alternative splicing results in multiple transcript variants.

## Recommended Dilutions

<b>WB</b>	1:1000 - 1:4000
<b>IF/ICC</b>	1:50 - 1:200
<b>IP</b>	1:20 - 1:50

## Immunogen Information

<b>Gene ID</b>	<b>Swiss Prot</b>
1654	O00571

**Immunogen**

Recombinant protein of human DDX3X

**Synonyms**

DBX; DDX3; HLP2; DDX14; CAP-Rf; MRX102; MRXSSB; DDX3X

## Contact

 | [www.abclonal.com](http://www.abclonal.com)

## Product Information


<b>Source</b>	<b>Isotype</b>	<b>Purification</b>
Mouse	IgG	Affinity purification

**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

# Validation Data



Western blot analysis of various lysates using DDX3X Mouse mAb (A10099) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 5s.