A11128

Leader in Biomolecular Solutions for Life Science

2 Publications

PTEN Mouse mAb

Catalog No.: A11128

Basic Information

Observed MW

Calculated MW

Polyclonal Antibody

Cross-Reactivity Human, Mouse, Rat

CloneNo number

Applications

54kDa

47kDa

Category

WB, ELISA

AMC0179



Background

This gene was identified as a tumor suppressor that is mutated in a large number of cancers at high frequency. The protein encoded by this gene is a phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase. It contains a tensin like domain as well as a catalytic domain similar to that of the dual specificity protein tyrosine phosphatases. Unlike most of the protein tyrosine phosphatases, this protein preferentially dephosphorylates phosphoinositide substrates. It negatively regulates intracellular levels of phosphatidylinositol-3,4,5-trisphosphate in cells and functions as a tumor suppressor by negatively regulating AKT/PKB signaling pathway. The use of a non-canonical (CUG) upstream initiation site produces a longer isoform that initiates translation with a leucine, and is thought to be preferentially associated with the mitochondrial inner membrane. This longer isoform may help regulate energy metabolism in the mitochondria. A pseudogene of this gene is found on chromosome 9. Alternative splicing and the use of multiple translation start codons results in multiple transcript variants encoding different isoforms.

Recommended Dilutions

Immunogen Information

WB

1:500 - 1:1000

Gene ID 5728 Swiss Prot P60484

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 204-403 of human PTEN (P60484).

Synonyms

BZS; DEC; CWS1; GLM2; MHAM; TEP1; MMAC1; PTEN1; 10q23del; PTENbeta; PTEN

Contact

Product Information

S www.abc

www.abclonal.com

lsotype IgG1,kappa Purification Affinity purification


Storage

Source

Mouse

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using PTEN Mouse mAb (A11128) at 1:1000 dilution. Secondary anthogy:HRP Goat Anti-Mouse IgG (H+L)(AS003) at 1:10000 dilution. Lysates / proteins: 25 µg per lane. Blocking buffer 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s. 100kDa-75kDa-

