

A1162

Leader in Biomolecular Solutions for Life Science



[KO Validated] HNRNPA2B1 Rabbit pAb

Catalog No.: A1162

KO Validated

6 Publications

Basic Information

Observed MW

36-38kDa

Calculated MW

37kDa

Category

Polyclonal Antibody

Applications

WB, IHC-P, IF/ICC, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene belongs to the A/B subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has two repeats of quasi-RRM domains that bind to RNAs. This gene has been described to generate two alternatively spliced transcript variants which encode different isoforms.

Recommended Dilutions

WB	1:1000 - 1:5000
IHC-P	1:50 - 1:200
IF/ICC	1:100 - 1:500

Immunogen Information

Gene ID

3181

Swiss Prot

P22626

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-180 of human HNRNPA2B1 (NP_112533.1).

Synonyms

RNPA2; HNRPA2; HNRPB1; SNRPB1; HNRNPA2; HNRNPB1; IBMPFD2; HNRPA2B1; B1

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.


Validation Data




Western blot analysis of lysates from wild type (WT) and HNRNPA2B1 knockout (KO) 293T cells, using [KO Validated] HNRNPA2B1 Rabbit pAb (A1162) at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.




Western blot analysis of various lysates, using [KO Validated] HNRNPA2B1 Rabbit pAb (A1162) at 1:3500 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.



Immunohistochemistry analysis of HNRNPA2B1 in paraffin-embedded rat lung tissue using [KO Validated] HNRNPA2B1 Rabbit pAb (A1162) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of HNRNPA2B1 in paraffin-embedded human cervix tissue using [KO Validated] HNRNPA2B1 Rabbit pAb (A1162) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of HNRNPA2B1 in paraffin-embedded human brain tissue using [KO Validated] HNRNPA2B1 Rabbit pAb (A1162) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.

Validation Data



Confocal immunofluorescence analysis of HeLa cells using [KO Validated] HNRNPA2B1 Rabbit pAb (A1162) at dilution of 1:400. Blue: DAPI for nuclear staining.