Leader in Biomolecular Solutions for Life Science

# **SIRT1 Mouse mAb**

Catalog No.: A11866



## **Basic Information**

### **Observed MW**

120kDa

### **Calculated MW**

82kDa

#### Category

SMab Recombinant Monoclonal Antibody

#### **Applications**

WB,IF/ICC,IP,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat, Monkey

# **Background**

This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Alternative splicing results in multiple transcript variants.

# **Recommended Dilutions**

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:100

1:20 - 1:50

# **Immunogen Information**

**Gene ID Swiss Prot** 23411 996EB6

### **Immunogen**

Recombinant protein of human SIRT1

## **Synonyms**

SIR2; SIR2L1; SIR2alpha; SIRT1

### **Contact**

ΙP

www.abclonal.com

## **Product Information**


SourceIsotypePurificationMouseIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from MEF(WT) cells, using SIRT1 Mouse mAb (A11866). Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.