Leader in Biomolecular Solutions for Life Science

VWC2L Rabbit pAb

Catalog No.: A17853



Basic Information

Observed MW

23kDa

Calculated MW

Category

Polyclonal Antibody

Applications

WB,ELISA

Cross-Reactivity

Human, Mouse

Background

Acts upstream of or within negative regulation of BMP signaling pathway and positive regulation of neuron differentiation. Located in extracellular space. Part of AMPA glutamate receptor complex. Is expressed in brain; central nervous system; hippocampus; hypothalamus; and thalamus. Orthologous to human VWC2L (von Willebrand factor C domain containing 2 like).

Recommended Dilutions

WB

1:500 - 1:2000

Immunogen Information

Gene ID 320460

Swiss Prot

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-50 of mouse VWC2L (NP 796138.2).

Synonyms

Brl; A830006F12Rik; VWC2L

Contact

•

www.abclonal.com

Product Information


Source Rabbit **Isotype** IgG **Purification**Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using VWC2L Rabbit pAb (A17853) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.