Leader in Biomolecular Solutions for Life Science

# **GGCX Rabbit pAb**

Catalog No.: A1806 1 Publications



### **Basic Information**

### **Observed MW**

110kDa

#### **Calculated MW**

88kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse

### **Background**

This gene encodes an integral membrane protein of the rough endoplasmic reticulum that carboxylates glutamate residues of vitamin K-dependent proteins to gamma carboxyl glutamate, a modification that is required for their activity. The vitamin K-dependent protein substrates have a propeptide that binds the enzyme, with carbon dioxide, dioxide, and reduced vitamin K acting as co-substrates. Vitamin K-dependent proteins affect a number of physiologic processes including blood coagulation, prevention of vascular calcification, and inflammation. Allelic variants of this gene have been associated with pseudoxanthoma elasticum-like disorder with associated multiple coagulation factor deficiency. Alternative splicing results in multiple transcript variants.

### **Recommended Dilutions**

WB

1:500 - 1:2000

### **Immunogen Information**

Gene ID 2677 **Swiss Prot** 

P38435

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 479-758 of human GGCX (NP 000812.2).

### **Synonyms**

VKCFD1; GGCX

### **Contact**

•

www.abclonal.com

### **Product Information**

**Source** Rabbit **Isotype** IgG


**Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of extracts of various cell lines, using GGCX antibody (A1806) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.