A1831

Leader in Biomolecular Solutions for Life Science

[KO Validated] PTBP1 Rabbit pAb

Catalog No.: A1831

KO Validated 1 Publications



Basic Information

Observed MW 57kDa

Calculated MW 60kDa

Category Polyclonal Antibody

Applications WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity Human, Mouse, Rat

Background

This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA-binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has four repeats of quasi-RNA recognition motif (RRM) domains that bind RNAs. This protein binds to the intronic polypyrimidine tracts that requires pre-mRNA splicing and acts via the protein degradation ubiquitin-proteasome pathway. It may also promote the binding of U2 snRNP to pre-mRNAs. This protein is localized in the nucleoplasm and it is also detected in the perinucleolar structure. Alternatively spliced transcript variants encoding different isoforms have been described.

Recommended Dilutions

1:50 - 1:100

Immunogen Information

1:500 - 1:1000	Gene ID	Swiss Prot
	5725	P26599
1:50 - 1:100		

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-290 of human PTBP1 (NP 114368.1).

Synonyms

PTB; PTB2; PTB3; PTB4; pPTB; HNRPI; PTB-1; PTB-T; HNRNPI; HNRNP-I; P1

Ð

WB

IHC-P

IF/ICC

Product Information

www.abclonal.com

Isotype lgG

Purification Affinity purification


Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts from wild type(WT) and PTBP1 knockout (KO) HeLa(KO) cells, using PTBP1 antibody: (Å1831) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lystes/proteins: 25µg per lane. Billotking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s. 70kDa-55kDa--PTBP1




HeLa

40kDa

35kDa-

B-actir

Immunohistochemistry analysis of paraffinembedded human colon carcinoma using [KO Validated] PTBP1 Rabbit pAb (A1831) at dilution of 1:25 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Confocal immunofluorescence analysis of U-2OS cells using PTBP1 antibody (A1831) at dilution of 1:50. Blue: DAPI for nuclear staining.