

A19585

Leader in Biomolecular Solutions for Life Science



[KO Validated] p53 Rabbit mAb

Catalog No.: A19585

KO Validated

Recombinant

10 Publications

Basic Information

Observed MW

53kDa/

Calculated MW

44kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IP,ELISA

Cross-Reactivity

Human,Rat

CloneNo number

ARC56085

Background

This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons from identical transcript variants (PMIDs: 12032546, 20937277).

Recommended Dilutions

WB 1:500 - 1:1000

IP 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells

Immunogen Information

Gene ID

7157

Swiss Prot

P04637

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-393 of human p53 (NP_000537.3).

Synonyms

P53; BCC7; LFS1; BMFS5; TRP53; 53

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from 293T cells using [KO Validated] p53 Rabbit mAb (A19585) at 1:1000 dilution .
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L)(AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020)
Exposure time: 1s.

Immunoprecipitation analysis of 300 µg extracts from 293T cells using 3 µg [KO Validated] p53 Rabbit mAb (A19585). Western blot was performed from the immunoprecipitate using [KO Validated] p53 Rabbit mAb (A19585) at a dilution of 1:1000.

