

A2216

Leader in Biomolecular Solutions for Life Science



TSG101/VPS23 Rabbit pAb

Catalog No.: A2216

19 Publications

Basic Information

Observed MW

44kDa

Calculated MW

44kDa

Category

Polyclonal Antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

The protein encoded by this gene belongs to a group of apparently inactive homologs of ubiquitin-conjugating enzymes. The gene product contains a coiled-coil domain that interacts with stathmin, a cytosolic phosphoprotein implicated in tumorigenesis. The protein may play a role in cell growth and differentiation and act as a negative growth regulator. In vitro steady-state expression of this tumor susceptibility gene appears to be important for maintenance of genomic stability and cell cycle regulation. Mutations and alternative splicing in this gene occur in high frequency in breast cancer and suggest that defects occur during breast cancer tumorigenesis and/or progression.

Recommended Dilutions

WB 1:500 - 1:1000

IF/ICC 1:50-1:200

Immunogen Information

Gene ID

7251

Swiss Prot

Q99816

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 291-390 of human TSG101/VPS23 (NP_006283.1).

Synonyms

TSG10; VPS23; TSG101/VPS23

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.


Validation Data




Western blot analysis of K-562, using TSG101/VPS23 Rabbit pAb (A2216) at 1:800 dilution.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.
 Lysates/proteins: 25ug per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 30s.



Western blot analysis of extracts from normal (control) and TSG101/VPS23 knockdown (KD) SH-SY5Y(KD) cells, using TSG101/VPS23 Rabbit pAb (A2216) at 1:800 dilution.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.
 Lysates/proteins: 25ug per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 30s.



Immunofluorescence analysis of Jurkat cells using TSG101/VPS23 Rabbit pAb (A2216) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using TSG101/VPS23 Rabbit pAb (A2216) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.