

A2368

Leader in Biomolecular Solutions for Life Science



## DiMethyl-Histone H3-K79 Rabbit pAb

Catalog No.: A2368

3 Publications

### Basic Information

#### Observed MW

17kDa

#### Calculated MW

16kDa

#### Category

Polyclonal Antibody

#### Applications

WB, IHC-P, IF/ICC, IP, ChIP, ChIP-seq, ELISA

#### Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

### Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

### Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:100
IF/ICC	1:50 - 1:200
IP	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells
ChIP	5µg antibody for 5µg-10µg of Chromatin
ChIP-seq	1:50 - 1:200

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Immunogen Information

#### Gene ID

8290/8350

#### Swiss Prot

Q16695/P68431

#### Immunogen

A synthetic dimethylated peptide around K79 of human Histone H3 (NP\_003520.1).

#### Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; DiMethyl-Histone H3-K79

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage



Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


Validation Data




Chromatin immunoprecipitations were performed with cross-linked chromatin from K-562 cells and DiMethyl-Histone H3-K79 Rabbit pAb (A2368). The ChIP sequencing results indicate the enrichment pattern of DiMethyl-Histone H3-K79 in selected genomic region and representative gene loci (GAPDH), as shown in figure.




Dot-blot analysis of all sorts of methylation peptides using DiMethyl-Histone H3-K79 antibody (A2368).




Immunohistochemistry analysis of DiMethyl-Histone H3-K79 in paraffin-embedded human mammary cancer using DiMethyl-Histone H3-K79 Rabbit pAb (A2368) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of DiMethyl-Histone H3-K79 in paraffin-embedded rat testis using DiMethyl-Histone H3-K79 Rabbit pAb (A2368) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of DiMethyl-




Immunofluorescence analysis of 293T cells

Validation Data

Histone H3-K79 in paraffin-embedded mouse testis using DiMethyl-Histone H3-K79 Rabbit pAb (A2368) at dilution of 1:200 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

using DiMethyl-Histone H3-K79 Rabbit pAb (A2368). Blue: DAPI for nuclear staining.



Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg DiMethyl-Histone H3-K79 antibody (A2368). Western blot was performed from the immunoprecipitate using DiMethyl-Histone H3-K79 antibody (A2368) at a dilution of 1:1000.

Chromatin Immunoprecipitation analysis of extracts of MCF7 cells, using DiMethyl-Histone H3-K79 antibody (A2368) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.