A2421

Leader in Biomolecular Solutions for Life Science

Asymmetric DiMethyl-Histone H3-R17 Rabbit pAb

ABclomal www.abclonal.com

Catalog No.: A2421 7 Publications

Basic Information

Observed MW 17kDa

Calculated MW 15kDa

Category Polyclonal Antibody

Applications WB, IF/ICC, ELISA

Cross-Reactivity Human, Mouse, Rat, Other (Wide Range Predicted)

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

Immunogen Information

WB	1:500 - 1:1000	Gene ID	Swiss Prot
		8290/8350	Q16695/P68431
IF/ICC	1:50 - 1:200		
		Immunogen	
		A synthetic peptide correspon	ding to a sequence within amino a

ino acids 1-100 of human histone H3 (NP 003520.1).

Synonyms

H3/j; H3C1; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FJ; H3C10; H3C11; HIST1H3J; Asymmetric DiMethyl-Histone H3-R17

Contact

Product Information

€

www.abclonal.com

Isotype lgG

Purification Affinity purification


Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using Asymmetric DiMethyl-Histone H3-R17 Rabbit pAb (A2421) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.

	H3R2		H3K4		H3R8		H3K9		H3R17		H3R26	
	1009	60n9	1009	50n9								
me0	0	0	0	0	0	0	0	0	0	0	0	0
me1	0	0	0	0	0	0	0	0	0	0	0	0
me2/ me2a	0	0	0	0	0	0	0	0	0	•	0	0
me3/ me2s	0	0	0	0	0	0	0	0	0	0	0	0
	H3K27		H3K36		H3K56		H3K79		H4R3		H4K20	
me0	0	0	0	0	0	0	0	0	0	0	0	0
me1	0	0	0	0	0	0	0	0	0	0	0	0
me2/ me2a	0	0	0	0	0	0	0	0	0	0	0	0
me3/ me2s	0	0	0	0	0	0	0	0.	0	0	0	0

Dot-blot analysis of all sorts of methylation peptides using Asymmetric DiMethyl-Histone H3-R17 antibody (A2421). Immunofluorescence analysis of 293T cells using Asymmetric DiMethyl-Histone H3-R17 Rabbit pAb (A2421). Blue: DAPI for nuclear staining.

H3R17me2a

DAP

Immunofluorescence analysis of HeLa cells using Asymmetric DiMethyl-Histone H3-R17

using Asymmetric DiMethyl-Histone H3-R17 Rabbit pAb (A2421) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.