Leader in Biomolecular Solutions for Life Science

# Nanog Rabbit pAb

Catalog No.: A3232 13 Publications



### **Basic Information**

### **Observed MW**

42kDa

#### **Calculated MW**

35kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,IHC-P,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

The protein encoded by this gene is a DNA binding homeobox transcription factor involved in embryonic stem (ES) cell proliferation, renewal, and pluripotency. The encoded protein can block ES cell differentiation and can also autorepress its own expression in differentiating cells. Two transcript variants encoding different isoforms have been found for this gene.

### **Recommended Dilutions**

**WB** 1:1000 - 1:2000

IHC-P 1:50 - 1:200

### **Immunogen Information**

**Gene ID**Swiss Prot
79923
Q9H9S0

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-105 of human Nanog (NP\_079141.2).

### **Synonyms**

NANOG; Nanog

### **Contact**

www.abclonal.com


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.




Western blot analysis of Rat brain, using Nanog Rabbit pAb (A3232) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.



Immunohistochemistry analysis of paraffinembedded Human testicular seminoma using Nanog Rabbit pAb (A3232) at dilution of 1:20 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.