Leader in Biomolecular Solutions for Life Science

# Asymmetric DiMethyl-Histone H4-R3 Rabbit pAb



Catalog No.: A7261

## **Basic Information**

## **Observed MW**

14kDa

## **Calculated MW**

11kDa

### Category

Mouse Monoclonal Antibody

## **Applications**

WB,IHC-P,ELISA

## **Cross-Reactivity**

Human, Mouse, Rat, Other (Wide Range Predicted)

# **Background**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in a histone cluster on chromosome 1. This gene is one of four histone genes in the cluster that are duplicated; this record represents the centromeric copy.

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

IHC-P 1:50 - 1:200

# **Immunogen Information**

Gene ID Swiss Prot 8359 P62805

### **Immunogen**

A synthetic asymmetric dimethylated peptide around R3 of human HIST2H4A (NP\_003529.1).

#### Synonyme

H4; H4/n; H4C1; H4C2; H4C3; H4C4; H4C5; H4C6; H4C8; H4C9; H4F2; H4FN; F0108; H4-16; H4C11; H4C12; H4C13; H4C15; H4C16; HIST2H4; HIST2H4A; Asymmetric DiMethyl-Histone H4-R3

## **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**




Western blot analysis of lysates from HeLa cells, using Asymmetric DiMethyl-Histone H4-R3 Rabbit pAb (A7261) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 300s.



Dot-blot analysis of all sorts of methylation peptides using Asymmetric DiMethyl-Histone H4-R3 antibody (A7261) at 1:1000 dilution.