Leader in Biomolecular Solutions for Life Science

# **TBL1XR1** Mouse mAb

Catalog No.: A9922



### **Basic Information**

### **Observed MW**

55kDa

### **Calculated MW**

56kDa

### Category

SMab Recombinant Monoclonal Antibody

#### **Applications**

WB,IHC-P,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse

## **Background**

This gene is a member of the WD40 repeat-containing gene family and shares sequence similarity with transducin (beta)-like 1X-linked (TBL1X). The protein encoded by this gene is thought to be a component of both nuclear receptor corepressor (N-CoR) and histone deacetylase 3 (HDAC 3) complexes, and is required for transcriptional activation by a variety of transcription factors. Mutations in these gene have been associated with some autism spectrum disorders, and one finding suggests that haploinsufficiency of this gene may be a cause of intellectual disability with dysmorphism. Mutations in this gene as well as recurrent translocations involving this gene have also been observed in some tumors.

## **Recommended Dilutions**

WB	1:500 - 1:2000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

## **Immunogen Information**

 Gene ID
 Swiss Prot

 79718
 Q9BZK7

### **Immunogen**

Recombinant protein of human TBL1XR1

## **Synonyms**

C21; DC42; IRA1; MRD41; TBLR1; TBL1XR1

### **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationMouseIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates using TBL1XR1 Mouse mAb (A9922) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Mouse  $\lg G$  (H+L) (AS003) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 20s.