Leader in Biomolecular Solutions for Life Science

MT-ND4 Rabbit pAb

Catalog No.: A9941 10 Publications



Basic Information

Observed MW

50kDa

Calculated MW

51kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Predicted to enable NADH dehydrogenase (ubiquinone) activity and ubiquinone binding activity. Predicted to contribute to NADH dehydrogenase activity. Predicted to be involved in several processes, including electron transport coupled proton transport; mitochondrial electron transport, NADH to ubiquinone; and mitochondrial respiratory chain complex I assembly. Located in mitochondrion. Human ortholog(s) of this gene implicated in Leber hereditary optic neuropathy; Parkinson's disease; macular degeneration; and schizophrenia. Orthologous to human MT-ND4 (mitochondrially encoded NADH:ubiquinone oxidoreductase core subunit 4).

Recommended Dilutions

WB

1:500 - 1:1000

Immunogen Information

Gene ID 17719

Swiss Prot

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 360-459 of mouse MT-ND4 (NP 904337.1).

Synonyms

ND4; MT-ND4

Contact

•

www.abclonal.com

Product Information


Source Rabbit **Isotype** IgG **Purification**Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using MT-ND4 antibody (A9941) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.