

α -Tubulin Rabbit pAb

Catalog No.: AC014

Basic Information

Observed MW

55kDa

Calculated MW

50kDa

Category

Loading control antibody

Applications

WB, IHC-P, IF/ICC

Cross-Reactivity

Human, Mouse, Rat

Background

Microtubules of the eukaryotic cytoskeleton perform essential and diverse functions and are composed of a heterodimer of alpha and beta tubulin. The genes encoding these microtubule constituents are part of the tubulin superfamily, which is composed of six distinct families. Genes from the alpha, beta and gamma tubulin families are found in all eukaryotes. The alpha and beta tubulins represent the major components of microtubules, while gamma tubulin plays a critical role in the nucleation of microtubule assembly. There are multiple alpha and beta tubulin genes and they are highly conserved among and between species. This gene encodes an alpha tubulin that is a highly conserved homolog of a rat testis-specific alpha tubulin. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

| | |
|---------------|----------------|
| WB | 1:500 - 1:2000 |
| IHC-P | 1:25 - 1:50 |
| IF/ICC | 1:50 - 1:200 |

Immunogen Information

Gene ID

7277

Swiss Prot

P68366

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-180 of human α -Tubulin (P68366).

SynonymsALS22; TUBA1; H2-ALPHA; α -Tubulin

Contact

 | www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of various lysates using α -Tubulin Rabbit pAb (AC014) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).




Immunohistochemistry analysis of α -Tubulin in paraffin-embedded rat brain using α -Tubulin Rabbit pAb (AC014) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




Immunohistochemistry analysis of α -Tubulin in paraffin-embedded rat kidney using α -Tubulin Rabbit pAb (AC014) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




Immunohistochemistry analysis of α -Tubulin in paraffin-embedded rat heart using α -Tubulin Rabbit pAb (AC014) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of C6 cells using α -Tubulin Rabbit pAb (AC014) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using α -Tubulin Rabbit pAb (AC014) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using α -Tubulin Rabbit pAb (AC014) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.