AC038

Leader in Biomolecular Solutions for Life Science

Recombinant

β-Actin Rabbit mAb

Catalog No.: AC038

126 Publications

Basic Information

Observed MW 45kDa

Calculated MW 42kDa

Category SMab Recombinant Monoclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human, Mouse, Rat, Chicken, Zebrafish, Pig

CloneNo number ARC5115-01

Background

This gene encodes one of six different actin proteins. Actins are highly conserved proteins that are involved in cell motility, structure, integrity, and intercellular signaling. The encoded protein is a major constituent of the contractile apparatus and one of the two nonmuscle cytoskeletal actins that are ubiquitously expressed. Mutations in this gene cause Baraitser-Winter syndrome 1, which is characterized by intellectual disability with a distinctive facial appearance in human patients. Numerous pseudogenes of this gene have been identified throughout the human genome.

Recommended Dilutions

Immunogen Information

WB

1:10000 - 1:100000

Gene ID

Swiss Prot P60709

 $\label{eq:starses} \begin{array}{l} \textbf{Immunogen} \\ \textbf{Recombinant protein of human } \beta \textbf{-Actin} \end{array}$

Synonyms BRWS1; PS1TP5BP1; β-Actin

Contact

€

Product Information

www.abclonal.com

lsotype IgG Purification Affinity purification

Storage


Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.



Validation Data



Western blot analysis of various lysates using β -Actin Rabbit mAb (AC038) at 1:50000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 60s.

Western blot analysis of dysates from HeLa cells, using β-Actin Rabbit mAb (AC038) at 1:10000-1:320000 dilutions. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per Jane. Blocking buffer: 3% nonfat dry milk in TBST. 70kDa – Detection: ECL Basic Kit (RM00020). Exposure time: 180s. SokDa – 40kDa – 35kDa – 25kDa –