

AE025

Leader in Biomolecular Solutions for Life Science



HRP-conjugated Mouse anti HA-Tag mAb

Catalog No.: AE025

1 Publications

Basic Information

Observed MW

53kDa(CD95)/65kDa(CD2)/53kDa(SCRN1)
)/26KD(HA(N
端)/42kDa(WDR77)/45kDa(NR3C2)

Calculated MW

Category

Mouse Monoclonal Antibody

Applications

WB,ELISA

Cross-Reactivity

Species independent

CloneNo number

AMC0557

Conjugate

HRP

Recommended Dilutions

WB 1:500 - 1:5000

Background

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

Immunogen Information

Gene ID

Swiss Prot

Immunogen

A synthetic peptide corresponding to HA tag.

Synonyms

HA;HA tag;HA-tag

Contact



www.abclonal.com

Product Information

Source

Mouse

Isotype

IgG1,Kappa


Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS with 50% glycerol,pH7.3.

Validation Data




Western blot analysis of extracts of normal 293T cells, 293T transfected with SCR1 Protein, using HRP-conjugated Mouse anti HA-Tag mAb (AE025) at 1:5000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.

Western blot analysis of extracts of normal 293T and 293F cells, 293T transfected with CD2 Protein and HA-N Protein, 293F transfected with CD95-N Protein using HRP-conjugated Mouse anti HA-Tag mAb (AE025) at 1:5000 dilution.

Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 35s.



Western blot analysis of extracts of normal 293T and 293F cells, 293T transfected with NR3C2-C Protein, 293F transfected with WDR77-C Protein using HRP-conjugated Mouse anti HA-Tag mAb (AE025) at 1:5000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.

