

AP0707

Leader in Biomolecular Solutions for Life Science



## Phospho-IkB $\alpha$ -S32 Rabbit mAb

Catalog No.: AP0707

Recombinant

70 Publications

### Basic Information

#### Observed MW

39kDa

#### Calculated MW

36kDa

#### Category

SMab Recombinant Monoclonal Antibody

#### Applications

WB, ELISA

#### Cross-Reactivity

Human, Mouse, Rat

#### CloneNo number

ARC0147

### Background

This gene encodes a member of the NF-kappa-B inhibitor family, which contain multiple ankrin repeat domains. The encoded protein interacts with REL dimers to inhibit NF-kappa-B/REL complexes which are involved in inflammatory responses. The encoded protein moves between the cytoplasm and the nucleus via a nuclear localization signal and CRM1-mediated nuclear export. Mutations in this gene have been found in ectodermal dysplasia anhidrotic with T-cell immunodeficiency autosomal dominant disease.

### Recommended Dilutions

WB 1:500 - 1:2000

### Immunogen Information

#### Gene ID

4792

#### Swiss Prot

P25963

#### Immunogen

A synthetic phosphorylated peptide around S32 of human IKB alpha (P25963).

#### Synonyms

IKBA; MAD-3; NFKBI; EDAID2; Phospho-IkB $\alpha$ -S32

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

## Validation Data



Western blot analysis of lysates from NIH/3T3 cells, using Phospho-IκBα-S32 Rabbit mAb (AP0707) at 1:1000 dilution. NIH/3T3 and C6 cells were treated by TNF-α (20 ng/mL) at 37°C for 30 minutes. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25μg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 120s.