

AP1385

Leader in Biomolecular Solutions for Life Science



Phospho-VEGF Receptor 2-Y1175 Rabbit mAb

Catalog No.: AP1385

Recombinant

Basic Information

Observed MW

230kDa

Calculated MW

152kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human

CloneNo number

ARC57293

Background

Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encodes one of the two receptors of the VEGF. This receptor, known as kinase insert domain receptor, is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors, including Rab GTPase, P2Y purine nucleotide receptor, integrin alphaVbeta3, T-cell protein tyrosine phosphatase, etc.. Mutations of this gene are implicated in infantile capillary hemangiomas.

Recommended Dilutions

WB 1:500 - 1:1000

Immunogen Information

Gene ID

3791

Swiss Prot

P35968

Immunogen

A synthetic phosphorylated peptide around Y1175 of human VEGF Receptor 2.

Synonyms

FLK1; CD309; VEGFR; VEGFR2; Phospho-VEGF Receptor 2-Y1175

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of A-431 lysates, using Phospho-VEGF Receptor 2-Y1175 antibody (AP1385) at 1:3000 dilution. A-431 cells were treated by EGF (100 ng/ml) at 37°C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.