Leader in Biomolecular Solutions for Life Science

ABclonal www.abclonal.com

HRP-conjugated Rabbit anti-Chicken IgY (IgG) (H+L)

Catalog No.: AS030 2 Publications

Basic Information

Observed MW

22-30kDa(Light chain),60-70kDa(Heavy chain)/30kDa(Light chain),70kDa(Heavy chain)

Calculated MW

Category

Secondary Antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Conjugate

HRP

Background

Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies . Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

Recommended Dilutions

ELISA 1:10000 - 1:200000

WB 1:10000 - 1:50000

IHC-P 1:500 - 1:5000

Immunogen Information

Gene ID Swiss Prot

Immunogen

Chicken IgY (IgG)

Synonyms

Contact

www.abclonal.com

Product Information


Source Isotype Purification
Rabbit Horseradish peroxidase conjugated IgG

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from Chicken $\lg Y$ using HRP Rabbit Anti-Chicken $\lg Y$ ($\lg G$) (H+L) (AS030) at 1:5000 dilution.

Lysates/proteins: 50ng - 100ng per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.

Western blot analysis of lysates from Chicken IgY using HRP Rabbit Anti-Chicken IgY (IgG) (H+L) (AS030) at 1:5000 dilution.

Lysates/proteins: 100ng per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

