

AS040

Leader in Biomolecular Solutions for Life Science



Rhodamine (TRITC)-conjugated Goat anti-Rabbit IgG (H+L)

Catalog No.: AS040

21 Publications

Basic Information

Observed MW

Calculated MW

Category

Secondary Antibody

Applications

IHC-P,IF/ICC,FC

Cross-Reactivity

Conjugate

Rhodamine. Ex:550nm. Em:570nm.

Background

Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies. Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

Recommended Dilutions

IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200
FC	1:50 - 1:200

Immunogen Information

Gene ID Swiss Prot

Immunogen

Rabbit IgG

Synonyms

Contact



www.abclonal.com

Product Information

Source

Goat

Isotype

TRITC conjugated IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.025% Sodium Azide,0.75% BSA,50% glycerol,pH7.3.


Validation Data



Immunofluorescence analysis of HeLa cells, using Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) (A3716) as the primary antibody at dilution of 1:100. The cells were incubated with the primary antibody overnight at 4°C.
Secondary antibody: Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) (AS040) at 1:200 dilution.
Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells, using Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) (A0942) as the primary antibody at dilution of 1:100. The cells were incubated with the primary antibody overnight at 4°C.
Secondary antibody: Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) (AS040) at 1:100 dilution.
Blue: DAPI for nuclear staining.



Flow cytometry: 1×10^6 K-562 cells (negative control, left) and A-431 cells (right) were surface-stained with Purified Rabbit anti-Human E-Cadherin mAb (5 μ l/Test, orange line) or secondary antibody only (blue line). Non-fluorescently stained K-562 and A-431 cells were used as blank control (red line). Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) (AS040, 1:200) was used as a secondary antibody.