

RP01150

Leader in Biomolecular Solutions for Life Science



Recombinant Human VEGF-A/VEGF165 Protein

Catalog No.: RP01150

Recombinant

2 Publications

Sequence Information

Species	Gene ID	Swiss Prot
Human	7422	P15692-4

Tags

N-His

Synonyms

VEGFA; MVCD1; VEGF; VPF; vascular endothelial growth factor
A;MVCD1;VEGF;VPF;L VEGFA;VEGF A

Product Information

Source	Purification
HEK293 cells	> 97% by SDS-PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Contact



www.abclonal.com

Background

Basic Information

Description

Recombinant Human VEGF-A/VEGF165 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala27-Arg191) of human VEGF165 (Accession #NP_001165097.1) fused with a 6×His tag at the N-terminus.


Bio-Activity

1. Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human VEGF165 at 1 μg/mL (100 μL/well) can bind Recombinant Human VEGFR2 with a linear range of 8-20 ng/mL. 2. Measured by its binding ability in a functional ELISA. Immobilized Human VEGF165 at 2 μg/mL (100 μL/well) can bind Human KDR with a linear range of 0.2-11.6 ng/mL.


Storage

Store the lyophilized protein at -20°C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.


Validation Data




Recombinant Human VEGF-A/VEGF165 Protein was determined by SDS-PAGE with Coomassie Blue, showing bands at 22 kDa and 27 kDa..




The purity of human VEGF165 Protein (Cat.RP01150) was greater than 90% as determined by SEC-HPLC.



Immobilized Recombinant Human VEGF165 at 1 µg/mL (100 µL/well) can bind Recombinant Human VEGFR2 with a linear range of 8-20 ng/mL.



Immobilized Recombinant Human VEGF165 at 2 µg/mL (100 µL/well) can bind Human KDR with a linear range of 0.2-11.6 ng/mL.



Recombinant Human VEGF165 stimulates cell proliferation of the human umbilical vein endothelial cells (HUVEC). The ED₅₀ for this effect is typically 0.19-0.78 ng/mL, corresponding to a specific activity of 1.28 × 10⁶~5.26 × 10⁶ units/mg.