Leader in Biomolecular Solutions for Life Science



# Recombinant Human B7-H1/PD-L1/CD274 Protein

Catalog No.: RP02045 Recombinant

### **Sequence Information**

Species	Gene ID	Swiss Prot
Human	29126	Q9NZQ7

Tags C-mFc

### Synonyms

B7-H; B7H1; PDL1; PD-L1; hPD-L1; PDCD1L1; PDCD1LG1;CD274;PDL1;B7H1;PD-L1;PDCD1L1;PDCD1LG1; B7-H; CD274 molecule

### **Product Information**

Source	Purification
HEK293 cells	> 95% by SDS-
	PAGE.

### Endotoxin

< 0.1 EU/µg of the protein by LAL method.

### Formulation

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

# Contact

€

www.abclonal.com

### Background

B7-H1, also known as PD-L1 and CD274, is an approximately 65 kDa transmembrane glycoprotein in the B7 family of immune regulatory molecules. PD-L1 has been identified as the ligand for the immunoinhibitory receptor programmed death-1 (PD1/PDCD1) and has been demonstrated to play a role in the regulation of immune responses and peripheral tolerance.

# **Basic Information**

### Description

Recombinant Human CD30 Ligand/TNFSF8 Protein is produced by mammalian expression system. The target protein is expressed with sequence (Phe19-Arg238) of human PD-L1 (Accession #Q9NZQ7) fused with an Fc tag at the C-terminus.


### **Bio-Activity**

Immobilized Human PD-L1 at 0.2  $\mu$ g/mL (100  $\mu$ L/well), dose response curve for Anti-PD-L1 Ab with the EC<sub>50</sub> of 1.46 ng/mL determined by ELISA.


### Storage

Store the lyophilized protein at -20°C to -80 °C for long term.<br/>br>After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.


# Validation Data




Human PD-L1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



Immobilized Human PD-L1, mFc Tag at 2µg/ml (100µl/Well) on the plate. Dose response curve for Human PD-1, hFc Tag with the  $EC_{s0}$  of 0.13µg/ml determined by ELISA.



The purity of Human PD-L1 is greater than 95% as determined by SEC-HPLC.



Immobilized Human PD-L1, mFc Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Anti-PD-L1 Antibody, hFc Tag with the EC<sub>50</sub> of 0.9µg/ml determined by ELISA.