Leader in Biomolecular Solutions for Life Science

Recombinant Human LRG1 Protein

www.abclonal.com

ABclonal

Catalog No.: RP02953

Recombinant

Sequence Information

Species Gene ID **Swiss Prot** Human 116844 P02750

Tags C-His

Synonyms LRG1;HMFT1766;LRG

Product Information

Source HEK293 cells **Purification** > 95% as

determined by HPLC

Endotoxin

<1EU/µg

Formulation

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

Diabetic nephropathy (DN) is an important public health concern of increasing proportions and the leading cause of end-stage renal disease (ESRD) in diabetic patients. It is one of the most common long-term microvascular complications of diabetes mellitus that is characterized by proteinuria and glomerular structural changes. LRG1 is a novel pro-angiogenic factors involved in the abnormal angiogenesis and renal fibrosis in DN.

Basic Information

Description

Recombinant Human LRG1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Val36-Gln347) of human LRG1 (Accession #NP 443204.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

Immobilized Human LRG1, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-LRG1 Antibody, hFc Tag with the EC₅₀ of 13.8ng/ml determined by ELISA.


Storage

Store the lyophilized protein at -20°C to -80°C for 12 months.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 $\,$

Avoid repeated freeze/thaw cycles.


Validation Data



Human LRG1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Human LRG1 is greater than 95% as determined by SEC-HPLC.



Immobilized Human LRG1, His Tag at $0.5\mu g/ml$ (100 $\mu l/Well$) on the plate. Dose response curve for Anti-LRG1 Antibody, hFc Tag with the EC $_{50}$ of 13.8ng/ml determined by ELISA.