Leader in Biomolecular Solutions for Life Science

# **APC Rabbit pAb**

Catalog No.: A0021



# **Basic Information**

### **Observed MW**

150kDa

#### **Calculated MW**

312kDa

#### Category

Mouse Monoclonal Antibody

## **Applications**

WB

#### **Cross-Reactivity**

Human

# **Background**

This gene encodes a tumor suppressor protein that acts as an antagonist of the Wnt signaling pathway. It is also involved in other processes including cell migration and adhesion, transcriptional activation, and apoptosis. Defects in this gene cause familial adenomatous polyposis (FAP), an autosomal dominant pre-malignant disease that usually progresses to malignancy. Mutations in the APC gene have been found to occur in most colorectal cancers, where disease-associated mutations tend to be clustered in a small region designated the mutation cluster region (MCR) and result in a truncated protein product.

# **Recommended Dilutions**

WB

1:500 - 1:1000

# **Immunogen Information**

Gene ID

324

Swiss Prot P25054

#### **Immunogen**

A synthetic peptide of human APC

# **Synonyms**

GS; DP2; DP3; BTPS2; DESMD; DP2.5; PPP1R46; APC

### **Contact**

•

www.abclonal.com

# **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from 293T cells, using APC Rabbit pAb (A0021). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.