

NQO1 Rabbit pAb

Catalog No.: A0047 **11 Publications**

Basic Information

Catalog No.

A0047

Observed MW

31kDa

Calculated MW

26kDa/27kDa/30kDa

Category

Primary antibody

Applications

WB,IF

Cross-Reactivity

Human, Mouse, Rat

Recommended Dilutions

WB	1:500 - 1:2000
IF	1:50 - 1:200

Contact

 | www.abclonal.com

Background

This gene is a member of the NAD(P)H dehydrogenase (quinone) family and encodes a cytoplasmic 2-electron reductase. This FAD-binding protein forms homodimers and reduces quinones to hydroquinones. This protein's enzymatic activity prevents the one electron reduction of quinones that results in the production of radical species. Mutations in this gene have been associated with tardive dyskinesia (TD), an increased risk of hematotoxicity after exposure to benzene, and susceptibility to various forms of cancer. Altered expression of this protein has been seen in many tumors and is also associated with Alzheimer's disease (AD). Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

Immunogen Information

Gene ID	Swiss Prot
1728	P15559

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-274 of human NQO1 (NP_000894.1).

Synonyms

DHQU;DIA4;DTD;NMOR1;NMORI;QR1;NQO1

Product Information


Source	Isotype	Purification
Rabbit	IgG	Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.


Validation Data




Western blot analysis of extracts of various cell lines, using NQO1 antibody (A0047) at 1:400 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10001 dilution. Lysates/proteins: 26ug per lane. Blocking buffer: 4% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00021). Exposure time: 3s.




Western blot analysis of extracts of Mouse liver, using NQO1 antibody (A0047) at 1:400 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10002 dilution. Lysates/proteins: 27ug per lane. Blocking buffer: 5% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00022). Exposure time: 30s.



Immunofluorescence analysis of NIH/3T3 cells using NQO1 Rabbit pAb (A0047) at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using NQO1 Rabbit pAb (A0047) at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using NQO1 Rabbit pAb (A0047) at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.