Leader in Biomolecular Solutions for Life Science

# **AKT2 Rabbit pAb**

Catalog No.: A0336 5 Publications



### **Basic Information**

# **Observed MW**

65kDa

### **Calculated MW**

56kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,IF/ICC,ELISA

#### **Cross-Reactivity**

Human, Mouse

# **Background**

This gene is a putative oncogene encoding a protein belonging to a subfamily of serine/threonine kinases containing SH2-like (Src homology 2-like) domains, which is involved in signaling pathways. The gene serves as an oncogene in the tumorigenesis of cancer cells For example, its overexpression contributes to the malignant phenotype of a subset of human ductal pancreatic cancers. The encoded protein is a general protein kinase capable of phophorylating several known proteins, and has also been implicated in insulin signaling.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:100

# **Immunogen Information**

Gene ID Swiss Prot 208 P31751

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 400-500 of human AKT2 (NP 001617.1).

### **Synonyms**

PKBB; PRKBB; HIHGHH; PKBBETA; RAC-BETA; AKT2

### **Contact**

www.abclonal.com

# **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts of various cell lines, using AKT2 antibody (A0336). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of HeLa cells using AKT2 antibody (A0336) at dilution of 1:100. Blue: DAPI for nuclear staining.