

A0416

Leader in Biomolecular Solutions for Life Science



# Activator protein 2 (AP-2/TFAP2A) Rabbit pAb

Catalog No.: A0416 **4 Publications**

## Basic Information

### Observed MW

50kDa

### Calculated MW

48kDa

### Category

Polyclonal Antibody

### Applications

WB,ELISA

### Cross-Reactivity

Human

## Background

The protein encoded by this gene is a transcription factor that binds the consensus sequence 5'-GCCNNNGGC-3'. The encoded protein functions as either a homodimer or as a heterodimer with similar family members. This protein activates the transcription of some genes while inhibiting the transcription of others. Defects in this gene are a cause of branchiooculofacial syndrome (BOFS). Three transcript variants encoding different isoforms have been found for this gene.

## Recommended Dilutions

WB 1:500 - 1:1000

## Immunogen Information

### Gene ID

7020

### Swiss Prot

P05549

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 120-200 of human Activator protein 2 (AP-2/TFAP2A) (NP\_003211.1).

### Synonyms

AP-2; BOFS; AP2TF; TFAP2; AP-2alpha; Activator protein 2 (AP-2/TFAP2A)

## Contact

 [www.abclonal.com](http://www.abclonal.com)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

## Validation Data

---



Western blot analysis of extracts of HeLa cells, using Activator protein 2 (AP-2/TFAP2A) antibody (A0416) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 180s.