ACADS Rabbit pAb

Catalog No.: A0945 3 Publications



Basic Information

Observed MW 44kDa

Calculated MW 44kDa

Category Polyclonal Antibody

Applications WB,IHC-P,ELISA

Cross-Reactivity Human, Mouse, Rat

Recommended Dilutions

Background

This gene encodes a tetrameric mitochondrial flavoprotein, which is a member of the acyl-CoA dehydrogenase family. This enzyme catalyzes the initial step of the mitochondrial fatty acid beta-oxidation pathway. Mutations in this gene have been associated with short-chain acyl-CoA dehydrogenase (SCAD) deficiency. Alternative splicing results in two variants which encode different isoforms.

Immunogen Information

WB	1:500 - 1:2000	Gene ID	Swiss Prot	
ІНС-Р	1:50 - 1:100	35	P16219	
	1.50 1.100	_		

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human ACADS (NP_000008.1).

Synonyms

SCAD; ACAD3; ACADS


Contact		Product Information		
€	www.abclonal.com	Source	Isotype	
-		Rabbit	IgG	

Purification Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data




Western blot analysis of various lysates using ACADS Rabbit pAb (A0945) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 15s.



Immunohistochemistry analysis of paraffin-embedded human liver cancer using ACADS Rabbit pAb (A0945) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human liver damage using ACADS Rabbit pAb (A0945) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.