Leader in Biomolecular Solutions for Life Science

[KO Validated] UBE2O Rabbit pAb

Catalog No.: A10036 KO Validated



Basic Information

Observed MW

200kDa

Calculated MW

141kDa

Category

Polyclonal Antibody

Applications

WB,IF/ICC,IP,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Enables ubiquitin conjugating enzyme activity and ubiquitin protein ligase activity. Involved in positive regulation of BMP signaling pathway; protein ubiquitination; and retrograde transport, endosome to Golgi. Located in cytoplasm and nuclear body.

Recommended Dilutions

WB 1:1000 - 1:4000

IF/ICC 1:50 - 1:200

IP 0.5μg-4μg antibody for 200μg-400μg extracts of

whole cells

Immunogen Information

Gene IDSwiss Prot
63893
Q9C0C9

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1073-1292 of human UBE2O (NP 071349.3).

Synonyms

E2-230K; 2O

Contact

www.abclonal.com


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates using [KO Validated] UBE2O Rabbit pAb (A10036) at 1:1000

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.


Western blot analysis of lysates from wild type (WT) and UBE2O knockout (KO) HeLa cells, using [KO Validated] UBE20 Rabbit pAb (A10036) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

309sates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).





Immunofluorescence analysis of HeLa cells using [KO Validated] UBE2O Rabbit pAb (A10036) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

Validation Data



Immunoprecipitation analysis of 200 μg extracts of A-431 cells using 3 μg UBE2O antibody (A10036). Western blot was performed from the immunoprecipitate using UBE2O antibody (A10036) at a dilution of 1:1000.