

ATP2B4 Rabbit pAb

Catalog No.: A10105

Basic Information

Catalog No.

A10105

Observed MW

138kDa

Calculated MW

124-137kDa

Category

Primary antibody

Applications

WB

Cross-Reactivity

Mouse

Recommended Dilutions

WB 1:1000 - 1:4000

Background

The protein encoded by this gene belongs to the family of P-type primary ion transport ATPases characterized by the formation of an aspartyl phosphate intermediate during the reaction cycle. These enzymes remove bivalent calcium ions from eukaryotic cells against very large concentration gradients and play a critical role in intracellular calcium homeostasis. The mammalian plasma membrane calcium ATPase isoforms are encoded by at least four separate genes and the diversity of these enzymes is further increased by alternative splicing of transcripts. The expression of different isoforms and splice variants is regulated in a developmental, tissue- and cell type-specific manner, suggesting that these pumps are functionally adapted to the physiological needs of particular cells and tissues. This gene encodes the plasma membrane calcium ATPase isoform 4. Alternatively spliced transcript variants encoding different isoforms have been identified.

Immunogen Information

Gene ID

493

Swiss Prot

P23634

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1141-1205 of human ATP2B4 (NP_001675.3).

Synonyms

ATP2B4;ATP2B2;MXRA1;PMCA4;PMCA4b;PMCA4x

Contact

 | www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of mouse brain, using ATP2B4 antibody (A10105) at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25ug per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 5s.