

# CACNA2D2 Rabbit pAb

Catalog No.: A10267 **1 Publications**

## Basic Information

**Catalog No.**

A10267

**Observed MW**

140-160kDa

**Calculated MW**

122kDa/129kDa/130kDa

**Category**

Primary antibody

**Applications**

WB

**Cross-Reactivity**

Mouse, Rat

## Recommended Dilutions

WB 1:200 - 1:2000

## Background

Calcium channels mediate the entry of calcium ions into the cell upon membrane polarization. This gene encodes the alpha-2/delta subunit of the voltage-dependent calcium channel complex. The complex consists of the main channel-forming subunit alpha-1, and auxiliary subunits alpha-2/delta, beta, and gamma. The auxiliary subunits function in the assembly and membrane localization of the complex, and modulate calcium currents and channel activation/inactivation kinetics. The subunit encoded by this gene undergoes post-translational cleavage to yield the extracellular alpha2 peptide and a membrane-anchored delta polypeptide. This subunit is a receptor for the antiepileptic drug, gabapentin. Mutations in this gene are associated with early infantile epileptic encephalopathy. Single nucleotide polymorphisms in this gene are correlated with increased sensitivity to opioid drugs. Alternative splicing results in multiple transcript variants encoding different isoforms.

## Immunogen Information

**Gene ID**

9254

**Swiss Prot**

Q9NY47

**Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 20-200 of human CACNA2D2 (NP\_001167522.1).

**Synonyms**

CACNA2D2;CACNA2D

## Contact

 | [www.abclonal.com](http://www.abclonal.com)

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**

Affinity purification


**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Validation Data

---



Western blot analysis of extracts of various cell lines, using CACNA2D2 antibody (A10267) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 1s.