

ASIC3 Rabbit pAb

Catalog No.: A10268

Basic Information

Catalog No.

A10268

Observed MW

59kDa

Calculated MW

44kDa/58kDa/60kDa

Category

Primary antibody

Applications

WB

Cross-Reactivity

Mouse, Rat

Recommended Dilutions

WB 1:500 - 1:2000

Background

This gene encodes a member of the degenerin/epithelial sodium channel (DEG/ENaC) superfamily. The members of this family are amiloride-sensitive sodium channels that contain intracellular N and C termini, two hydrophobic transmembrane regions, and a large extracellular loop, which has many cysteine residues with conserved spacing. The member encoded by this gene is an acid sensor and may play an important role in the detection of lasting pH changes. In addition, a heteromeric association between this member and acid-sensing (proton-gated) ion channel 2 has been observed as proton-gated channels sensitive to gadolinium. Alternatively spliced transcript variants have been described.

Immunogen Information

Gene ID

9311

Swiss Prot

Q9UHC3

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 62-320 of human ASIC3 (NP_064717.1).

Synonyms

ASIC3;ACCN3;DRASIC;SLNAC1;TNaC1

Contact

 | www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using ASIC3 antibody (A10268) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.