

ACE1 Rabbit mAb

Catalog No.: A11357 **Recombinant**

Basic Information

Observed MW

170kDa

Calculated MW

150kDa

Category

Primary antibody

Applications

ELISA,WB,IP

Cross-Reactivity

Mouse, Rat

CloneNo number

ARC0577

Recommended Dilutions

WB	1:500 - 1:1000
IP	0.5µg-4µg antibody for 400µg-600µg extracts of whole cells

Contact

 | www.abclonal.com

Background

This gene encodes an enzyme involved in blood pressure regulation and electrolyte balance. It catalyzes the conversion of angiotensin I into a physiologically active peptide angiotensin II. Angiotensin II is a potent vasopressor and aldosterone-stimulating peptide that controls blood pressure and fluid-electrolyte balance. This angiotensin converting enzyme (ACE) also inactivates the vasodilator protein, bradykinin. Accordingly, the encoded enzyme increases blood pressure and is a drug target of ACE inhibitors, which are often prescribed to reduce blood pressure. This enzyme additionally plays a role in fertility through its ability to cleave and release GPI-anchored membrane proteins in spermatozoa. Many studies have associated the presence or absence of a 287 bp Alu repeat element in this gene with the levels of circulating enzyme. This polymorphism, as well as mutations in this gene, have been implicated in a wide variety of diseases including cardiovascular pathophysiologies, psoriasis, renal disease, stroke, and Alzheimer's disease. Regulation of the homologous ACE2 gene may be involved in progression of disease caused by several human coronaviruses, including SARS-CoV and SARS-CoV-2. Alternative splicing results in multiple transcript variants encoding both somatic (sACE) and male-specific testicular (tACE) isoforms.

Immunogen Information

Gene ID	Swiss Prot
1636	P12821

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1200-1306 of human ACE1 (P12821).

Synonyms

DCP; ACE1; DCP1; CD143

Product Information


Source	Isotype	Purification
Rabbit	IgG	Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Immunoprecipitation analysis of 600 μ g extracts of Mouse kidney cells using 3 μ g ACE1 antibody (A11357). Western blot was performed from the immunoprecipitate using ACE1 antibody (A11357) at a dilution of 1:1000.



Western blot analysis of various lysates using ACE1 Rabbit mAb (A11357) at 1: 1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.