

A11399

Leader in Biomolecular Solutions for Life Science



δ-Catenin/p120 Catenin Rabbit mAb

Catalog No.: A11399 **Recombinant** **1 Publications**

Basic Information

Observed MW

100kDa/110kDa

Calculated MW

108kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IHC-P,IP,ELISA

Cross-Reactivity

Human,Mouse,Rat

CloneNo number

ARC0586

Background

This gene encodes a member of the Armadillo protein family, which function in adhesion between cells and signal transduction. Multiple translation initiation codons and alternative splicing result in many different isoforms being translated. Not all of the full-length natures of the described transcript variants have been determined. Read-through transcription also exists between this gene and the neighboring upstream thioredoxin-related transmembrane protein 2 (TMX2) gene.

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

IP 0.5µg-4µg antibody for
400µg-600µg extracts of
whole cells

Immunogen Information

Gene ID

1500

Swiss Prot

O60716

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 869-968 of human δ-Catenin/p120 Catenin (O60716).

Synonyms

CAS; p120; BCDS2; CTNND; P120CAS; P120CTN; p120(CAS); p120(CTN); δ-Catenin/p120 Catenin

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using δ -Catenin/p120 Catenin Rabbit mAb (A11399) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 μ g per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 3min.



Immunohistochemistry analysis of δ -Catenin/p120 Catenin in paraffin-embedded human breast using δ -Catenin/p120 Catenin Rabbit mAb (A11399) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of δ -Catenin/p120 Catenin in paraffin-embedded human placenta using δ -Catenin/p120 Catenin Rabbit mAb (A11399) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunoprecipitation analysis of 600 μ g extracts of Mouse testis cells using 3 μ g δ -Catenin/p120 Catenin antibody (A11399). Western blot was performed from the immunoprecipitate using δ -Catenin/p120 Catenin antibody (A11399) at a dilution of 1:1000.