

# [KO Validated] Bax Rabbit pAb

Catalog No.: A11550 **KO** **Validated** **1 Publications**

## Basic Information

### Catalog No.

A11550

### Observed MW

23kDa

### Calculated MW

4kDa/12kDa/15kDa/18kDa/19kDa/21kDa/24kDa

### Category

Primary antibody

### Applications

WB, IF, IP

### Cross-Reactivity

Human, Mouse, Rat

## Recommended Dilutions

**WB** 1:500 - 1:1000**IF** 1:50 - 1:100**IP** 1:50 - 1:200

## Contact

 | [www.abclonal.com](http://www.abclonal.com)

## Background

The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein forms a heterodimer with BCL2, and functions as an apoptotic activator. This protein is reported to interact with, and increase the opening of, the mitochondrial voltage-dependent anion channel (VDAC), which leads to the loss in membrane potential and the release of cytochrome c. The expression of this gene is regulated by the tumor suppressor P53 and has been shown to be involved in P53-mediated apoptosis. Multiple alternatively spliced transcript variants, which encode different isoforms, have been reported for this gene.

## Immunogen Information

### Gene ID

581

### Swiss Prot

Q07812

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-100 of human Bax (NP\_620116.1).

### Synonyms

BCL2L4;BAX

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


## Validation Data



Western blot analysis of extracts of various cell lines, using Bax antibody (A11550) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 60s.




Western blot analysis of extracts from normal (control) and Bax knockout (KO) 293T cells, using Bax antibody (A11550) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 5s.




Immunoprecipitation analysis of 200ug extracts of HeLa cells using 3ug Bax antibody (A11550). Western blot was performed from the immunoprecipitate using Bax antibody (A11550) at a dilution of 1:1000.

## Validation Data

---



Immunofluorescence analysis of C6 cells using Bax antibody (A11550) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using Bax antibody (A11550) at dilution of 1:100. Blue: DAPI for nuclear staining.