

A11625

Leader in Biomolecular Solutions for Life Science



Albumin Rabbit mAb

Catalog No.: A11625 **Recombinant** **1 Publications**

Basic Information

Observed MW

70kDa

Calculated MW

69kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,ELISA

Cross-Reactivity

Human,Mouse,Rat,Cow

CloneNo number

ARC0647

Background

Binds water, Ca²⁺, Na⁺, K⁺, fatty acids, hormones, bilirubin and drugs. Its main function is the regulation of the colloidal osmotic pressure of blood. Major zinc transporter in plasma, typically binds about 80% of all plasma zinc (By similarity. Major calcium and magnesium transporter in plasma, binds approximately 45% of circulating calcium and magnesium in plasma (Probable. Potentially has more than two calcium-binding sites and might additionally bind calcium in a non-specific manner. The shared binding site between zinc and calcium at residue Asp-272 suggests a crosstalk between zinc and calcium transport in the blood (Probable. The rank order of affinity is zinc > calcium > magnesium (Probable. Binds to the bacterial siderophore enterobactin and inhibits enterobactin-mediated iron uptake of E.coli, and may thereby limit the utilization of iron and growth of enteric bacteria such as E.coli. Does not prevent iron uptake by the bacterial siderophore aerobactin.

Recommended Dilutions

WB 1:2000 - 1:10000

Immunogen Information

Gene ID

280717

Swiss Prot

P02769

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 100-200 of bovine Serum Albumin / BSA (P02769).

Synonyms

Serum albumin; GIG20; GIG42; Albumin

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using Bovine Serum Albumin / BSA Rabbit mAb (A11625) at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 1s.