ADSS Rabbit pAb

Catalog No.: A12398



Basic Information

Observed MW 50kDa

Calculated MW 50kDa

Category Polyclonal Antibody

Applications WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity Human, Mouse

Recommended Dilutions

Background

This gene encodes the enzyme adenylosuccinate synthetase which catalyzes the first committed step in the conversion of inosine monophosphate to adenosine monophosphate. A pseudogene of this gene is found on chromosome 17.

Immunogen Information

WB	1:500 - 1:2000 1:50 - 1:100	Gene ID 159	Swiss Prot P30520	
IHC-P	1.50 - 1.100			
IF/ICC	1:50 - 1:100	Immunogen Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human ADSS (NP_001117.2).		

Synonyms


ADEH; ADSS; ADSS 2

Contact		Product Information			
•	www.abcional.com	Source Rabbit	Isotype IgG	Purification Affinity purification	

Storage


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data




Western blot analysis of various lysates using ADSS Rabbit pAb (A12398) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.




Immunohistochemistry analysis of paraffin-embedded human lung cancer using ADSS Rabbit pAb (A12398) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




Immunohistochemistry analysis of paraffin-embedded human colon using ADSS Rabbit pAb (A12398) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human esophageal cancer using ADSS Rabbit pAb (A12398) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse brain using ADSS Rabbit pAb (A12398) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of HeLa cells using ADSS Rabbit pAb (A12398). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.