Leader in Biomolecular Solutions for Life Science

# α-Synuclein Rabbit pAb

Catalog No.: A13354 2 Publications



# **Basic Information**

## **Observed MW**

18kDa

### **Calculated MW**

14kDa

### Category

Polyclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Alpha-synuclein is a member of the synuclein family, which also includes beta- and gamma-synuclein. Synucleins are abundantly expressed in the brain and alpha- and beta-synuclein inhibit phospholipase D2 selectively. SNCA may serve to integrate presynaptic signaling and membrane trafficking. Defects in SNCA have been implicated in the pathogenesis of Parkinson disease. SNCA peptides are a major component of amyloid plaques in the brains of patients with Alzheimer's disease. Alternatively spliced transcripts encoding different isoforms have been identified for this gene.

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

IF/ICC 1:50 - 1:200

# **Immunogen Information**

**Gene ID**Swiss Prot
6622
P37840

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 59-140 of human alpha-Synuclein (NP 000336.1).

### **Synonyms**

PD1; NACP; PARK1; PARK4; α-Synuclein

### **Contact**

www.abclonal.com


# **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.




Western blot analysis of extracts of various cell lines, using  $\alpha\textsc{-Synuclein}$  antibody (A13354) at 1:600

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Immunofluorescence analysis of mouse brain using α-Synuclein Rabbit pAb (A13354) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.