# ABclonal www.abclonal.com

## **AKR1C3 Rabbit pAb**

Catalog No.: A13568 2 Publications

## **Basic Information**

#### **Observed MW**

37kDa

#### **Calculated MW**

37kDa

#### Category

Primary antibody

## **Applications**

ELISA, WB, IHC-P

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes a member of the aldo/keto reductase superfamily, which consists of more than 40 known enzymes and proteins. These enzymes catalyze the conversion of aldehydes and ketones to their corresponding alcohols by utilizing NADH and/or NADPH as cofactors. The enzymes display overlapping but distinct substrate specificity. This enzyme catalyzes the reduction of prostaglandin (PG) D2, PGH2 and phenanthrenequinone (PQ), and the oxidation of 9alpha,11beta-PGF2 to PGD2. It may play an important role in the pathogenesis of allergic diseases such as asthma, and may also have a role in controlling cell growth and/or differentiation. This gene shares high sequence identity with three other gene members and is clustered with those three genes at chromosome 10p15-p14. Three transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

**WB** 1:1000 - 1:5000

**IHC-P** 1:50 - 1:200

## **Immunogen Information**

**Gene ID Swiss Prot**8644
P42330

## **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-323 of human AKR1C3 (NP\_003730.4).

### **Synonyms**

DD3; DDX; PGFS; HAKRB; HAKRe; HA1753; HSD17B5; hluPGFS; AKR1C3

#### **Contact**

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates, using AKR1C3 antibody (A13568) at 1:2000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded rat heart using AKR1C3 antibody (A13568) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.