# ABclonal www.abclonal.com

## **DNMT3L Rabbit pAb**

Catalog No.: A13591

## **Basic Information**

#### **Observed MW**

45kDa

#### **Calculated MW**

44kDa

#### Category

Primary antibody

## **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse

## **Background**

CpG methylation is an epigenetic modification that is important for embryonic development, imprinting, and X-chromosome inactivation. Studies in mice have demonstrated that DNA methylation is required for mammalian development. This gene encodes a nuclear protein with similarity to DNA methyltransferases, but is not thought to function as a DNA methyltransferase as it does not contain the amino acid residues necessary for methyltransferase activity. However, it does stimulate de novo methylation by DNA cytosine methyltransferase 3 alpha and is thought to be required for the establishment of maternal genomic imprints. This protein also mediates transcriptional repression through interaction with histone deacetylase 1. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

WB

1:500 - 1:2000

## **Immunogen Information**

Gene ID

**Swiss Prot** Q9UJW3

29947

## **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 151-386 of human DNMT3L (NP\_787063.1).

### **Synonyms**

DNMT3L

#### **Contact**

•

www.abclonal.com

## **Product Information**

**Source** Rabbit Isotype

IgG


**Purification** 

Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of extracts of various cell lines, using DNMT3L antibody (A13591) at 1:1000

dilutior

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.