# ABclonal® www.abclonal.com

## **ACSS2 Rabbit pAb**

Catalog No.: A13596

## **Basic Information**

## **Observed MW**

78kDa

## **Calculated MW**

79kDa

## Category

Primary antibody

#### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Rat

## **Background**

This gene encodes a cytosolic enzyme that catalyzes the activation of acetate for use in lipid synthesis and energy generation. The protein acts as a monomer and produces acetyl-CoA from acetate in a reaction that requires ATP. Expression of this gene is regulated by sterol regulatory element-binding proteins, transcription factors that activate genes required for the synthesis of cholesterol and unsaturated fatty acids. Alternative splicing results in multiple transcript variants.

## **Recommended Dilutions**

WB

1:500 - 1:1000

## **Immunogen Information**

**Gene ID** 55902

**Swiss Prot** 

Q9NR19

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 462-701 of human ACSS2 (NP\_061147.1).

#### **Synonyms**

ACS; ACSA; ACAS2; ACECS; AceCS1; dJ1161H23.1; ACSS2

## **Contact**

•

www.abclonal.com

## **Product Information**

Source

Isotype


**Purification** 

Rabbit IgG Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of extracts of various cell lines, using ACSS2 antibody (A13596) at 1:1000

dilution

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.