# **ABclonal** www.abclonal.com

## **CARS Rabbit pAb**

Catalog No.: A13676

## **Basic Information**

#### **Observed MW**

85kDa

## **Calculated MW**

85kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes a class 1 aminoacyl-tRNA synthetase, cysteinyl-tRNA synthetase. Each of the twenty aminoacyl-tRNA synthetases catalyzes the aminoacylation of a specific tRNA or tRNA isoaccepting family with the cognate amino acid. This gene is one of several located near the imprinted gene domain on chromosome 11p15.5, an important tumor-suppressor gene region. Alterations in this region have been associated with Beckwith-Wiedemann syndrome, Wilms tumor, rhabdomyosarcoma, adrenocortical carcinoma, and lung, ovarian and breast cancers. Alternative splicing of this gene results in multiple transcript variants.

## **Recommended Dilutions**

WB

1:500 - 1:2000

## **Immunogen Information**

**Gene ID** 

**Swiss Prot** P49589

833

**Immunogen** 

Recombinant fusion protein containing a sequence corresponding to amino acids 469-748 of human CARS (NP\_001742.1).

#### **Synonyms**

CARS; MDBH; CYSRS; MCDDBH; MGC:11246

#### **Contact**

•

www.abclonal.com

## **Product Information**

Source Rabbit

**Purification** 

Affinity purification


## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Isotype

IgG



Western blot analysis of various lysates using CARS Rabbit pAb (A13676) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.