Leader in Biomolecular Solutions for Life Science

TFAP2B Rabbit pAb

Catalog No.: A13885



Basic Information

Observed MW 45kDa

Calculated MW 50kDa

Category Primary antibody

Applications ELISA,WB

Cross-Reactivity Mouse

Background

This gene encodes a member of the AP-2 family of transcription factors. AP-2 proteins form homo- or hetero-dimers with other AP-2 family members and bind specific DNA sequences. They are thought to stimulate cell proliferation and suppress terminal differentiation of specific cell types during embryonic development. Specific AP-2 family members differ in their expression patterns and binding affinity for different promoters. This protein functions as both a transcriptional activator and repressor. Mutations in this gene result in autosomal dominant Char syndrome, suggesting that this gene functions in the differentiation of neural crest cell derivatives.

Recommended Dilutions

Immunogen Information

WB

Gene ID 7021

Swiss Prot Q92481

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-160 of human TFAP2B (NP_003212.2).

Synonyms

PDA2; AP-2B; AP2-B; AP-2beta; TFAP2B

Product Information


\odot	<u>www.abclonal.com</u>	Source
		Rabbit

1:500 - 1:2000

lsotype IgG Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of extracts of mouse skeletal muscle, using TFAP2B antibody (A13885) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.