# **ABclonal** www.abclonal.com

## **SLC22A2 Rabbit pAb**

Catalog No.: A14061 1 Publications

## **Basic Information**

#### **Observed MW**

95kDa

## **Calculated MW**

63kDa

#### Category

Primary antibody

## **Applications**

ELISA,WB

#### **Cross-Reactivity**

Mouse

## **Background**

Polyspecific organic cation transporters in the liver, kidney, intestine, and other organs are critical for elimination of many endogenous small organic cations as well as a wide array of drugs and environmental toxins. This gene is one of three similar cation transporter genes located in a cluster on chromosome 6. The encoded protein contains twelve putative transmembrane domains and is a plasma integral membrane protein. It is found primarily in the kidney, where it may mediate the first step in cation reabsorption.

## **Recommended Dilutions**

WB

1:500 - 1:2000

## **Immunogen Information**

**Gene ID** 6582

**Swiss Prot** 

015244

## **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 44-140 of human SLC22A2 (NP\_003049.2).

#### **Synonyms**

OCT2; SLC22A2

## **Contact**

•

www.abclonal.com

## **Product Information**

Source

Isotype

**Purification** 


Affinity purification

Rabbit IgG

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of extracts of mouse kidney, using SLC22A2 antibody (A14061) at 1:1000 ....

dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.